

Side Effects - Quiz 7

LaRue Medical Literacy Exercises

Drug Facts	
Active Ingredient (in each tablet) <i>Chlorpheniramine Maleate 2 mg</i>	Purpose <i>Antihistamine</i>
<i>Uses temporarily relieves these symptoms due to allergies</i> <div style="display: flex; justify-content: space-around;"> ■ <i>sneezing</i> ■ <i>runny nose</i> </div>	
Warnings <i>Ask a doctor before use if you have</i> <div style="display: flex; justify-content: space-around;"> ■ <i>glaucoma</i> ■ <i>a breathing problem such as emphysema</i> </div>	
When using this product <div style="list-style-type: none; padding-left: 0;"> <ul style="list-style-type: none"> ■ <i>drowsiness may occur</i> ■ <i>be careful when driving a motor vehicle or operating machinery</i> </div>	
Directions	
<i>Adults and children 12 and over</i>	<i>take 2 tablets every 4 to 6 hours not more than 12 tablets in 24 hours</i>
<i>Children 6 years to under 12 years</i>	<i>take 1 tablet every 4 to 6 hours not more than 6 tablets in 24 hour</i>
<i>Children under 6 years</i>	<i>ask a doctor</i>

1. If this medicine makes you drowsy, how do you feel ?

- A. hot
- B. tired
- C. cold
- D. hungry

2. What side effect makes it dangerous to drive?

- A. drowsiness
- B. glaucoma
- C. emphysema
- D. sneezing

3. People with this disease might have bad side effects from this medicine.

- A. people with allergies
- B. people with drowsiness
- C. people with runny noses
- D. people with emphysema

Drug Facts	
Active Ingredient (in each tablet)	Purpose
Pseudoephedrine HCL 30 mg.....	Nasal Decongestant
Uses temporarily relieves nasal congestion due to	
<ul style="list-style-type: none"> ■ common cold ■ respiratory allergies 	
Warnings	
Ask a doctor before use if you have	
<ul style="list-style-type: none"> ■ high blood pressure ■ heart disease ■ thyroid disease ■ diabetes 	
When using this product	
<ul style="list-style-type: none"> ■ do not use more than directed ■ Stop use and ask a doctor if you get nervous, dizzy or sleepless 	
Directions	
Adults and children 12 and over	take 2 tablets every 4 to 6 hours not more than 4 doses in 24 hours
Children 6 years to under 12 years	take 1 tablet every 4 to 6 hours not more than 4 doses in 24 hours
Children under 6 years	ask a doctor

4. What is not a side effect of this medicine?

- A. nervousness
- B. dizziness
- C. diabetes
- D. sleeplessness

5. If this medicine makes you sleepless...

- A. you sleep too much
- B. you are always sleepy
- C. you can't breathe
- D. you can't sleep

6. If this medicine makes you dizzy...

- A. it's hard to breathe
- B. it's hard to walk
- C. it's hard to sleep
- D. it's hard to see

7. People with this disease might have bad side effects from this medicine.

- A. heart disease
- B. kidney disease
- C. allergies
- D. cancer

Drug Facts	
Active Ingredient (in each tablet)	Purposes
Ibuprofen 200 mg.....	Pain Reliever/fever reducer
Uses temporarily relieves minor aches and pains due to <ul style="list-style-type: none"> ■ headache ■ muscular aches ■ backache 	
Warnings Ibuprofen may cause a severe allergic reaction which may be <ul style="list-style-type: none"> ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock 	
When using this product <ul style="list-style-type: none"> ■ take with food or milk if stomach upset occurs ■ Stop use and ask a doctor if pain gets worse or lasts more than 10 days 	
Directions	
Adults and children 12 and over	Take 1 tablet every 4 to 6 hours Do not exceed 6 tablets in 24 hours
Children under 12 years	ask a doctor

8. If you get hives as a side effect of this medicine, you have....

- A. a headache
- B. an upset stomach
- C. a skin rash
- D. wheezing

10. If you get facial swelling as a side effect of this medicine...

- A. your face gets red
- B. your face gets white
- C. your face gets hot
- D. your face gets bigger

9. If you get asthma as a side effect of this medicine...

- A. you can't walk
- B. you can't breathe
- C. you can't see
- D. you can't hear